

## **Fast Fulvic™**

Soil Balancing

Cropping

Pasture & Animal

Foliar & Fertigation

**An 8% fulvic acid liquid possessing a  
myriad of yield-enhancing qualities.**

### *Description*

**Fulvic Acid is a powerful organic extract from humates which offers a host of benefits for both the soil and the plant.**

### *Benefits*

- Australian Certified Organic (ACO) Registered Farm Input 456A1.
- Convenient liquid formulation.
- Natural chelating agent.
- Increases nutrient uptake.
- Promotes beneficial soil microbes.
- Improves availability of rock minerals.
- Increases crop yield and quality.
- Compatible with a wide range of other inputs.



### *Packaging*

1 L  
5 L  
20 L  
200 L  
1000 L



## Application Rates

### Foliar Spray:

#### Horticulture/Turf

0.5 L per 100 L water up to 3 L/ha equivalent  
Apply weekly or as required  
Include with all nutrient foliar sprays  
(confirm compatibility)

#### Broadacre Crops

1 L/ha in 60 – 100 L water, depending on canopy closure  
Include with all nutrient foliar sprays  
(confirm compatibility)

#### Pasture

1 – 2 L/ha in 60 – 100 L water Include with all nutrient foliar sprays (confirm compatibility)

#### Spot-Spraying

50 mL per 10 L water  
Apply as required

### Fertigation:

#### Horticulture/Turf

3 – 5 L/ha  
Apply weekly or as required Include with all nutrient applications (confirm compatibility)

### Home Garden:

Dilute 15 mL per 9 L watering can and apply the diluted mix at a rate of 1 litre per square metre  
Apply fortnightly or as required

## Instructions

**Shake or stir well before use.** Where higher water rates are required for good foliar coverage, do not exceed maximum product rate per hectare. Sensitive foliage or plants should be test sprayed prior to full scale application. Frequencies are a guideline only and can be adjusted according to farm practices and/or in conjunction with crop monitoring. Ensure suitable spray conditions (e.g. temperature, humidity, wind-speed), prior to foliar application. Compatibility and/or performance cannot be guaranteed when combining with other products (including various herbicide formulations). Jar test for compatibility and spray on a small area before widespread application.

### IMPORTANT

- Avoid spraying close to harvest if produce is susceptible to staining.
- **Note:** Product in 1000 L tanks may weigh over 1 tonne. Forklift with 2 tonne lifting capacity is recommended.

### STORAGE & HANDLING

- Store in a cool dry area out of direct sunlight. Do not pre-mix or store in diluted form. Seal lid/cap immediately after use.

## Typical Analysis W/V

Fulvic Acid .....	8%
SG .....	1.06
pH .....	3.5 – 4.5
Conductivity .....	20 – 40 mS/cm
Appearance .....	Brown liquid